

PAPER • OPEN ACCESS

## Preface

To cite this article: 2020 *IOP Conf. Ser.: Mater. Sci. Eng.* **852** 011001

View the [article online](#) for updates and enhancements.

## 2nd Tarumanagara International Conference on the Applications of Technology and Engineering 2019

### Preface

On behalf of the organizing committee of 2<sup>nd</sup> Tarumanagara International Conference on the Applications of Technology and Engineering (TICATE) 2019, I would like to welcome all delegates to Jakarta, Indonesia with great pleasure. Being held from November 21 to 22, 2019 at Campus I- Jl. Letjen. S. Parman No. 1, Jakarta, the international conference is organized by Universitas Tarumanagara (UNTAR) and technically sponsored by IOP Publisher.

TICATE 2019 has attracted many academicians, scientists, engineers, postgraduates and other professionals from many countries. This conference accepted 215 papers from 7 different countries, those are Australia, Taiwan, India, Malaysia, Japan, Peru and Indonesia. The aim of the conference is to promote exchange of ideas among engineers, researchers, and scientists active in the related areas of technology and engineering.

Our special thank goes to our Rector, Prof. Dr. Agustinus Purna Irawan, who has initiate this international conference, to our Plenary Speakers, Dr.-Ing. Joewono Prasetijo from Universiti Tun Hussein Onn, Malaysia, Prof. Dr. Tjokorda Gde Tirta Nindhia from Udayana University, Indonesia, Prof. Dr. Srikantappa A.S. from Cauvery Institute of Technology, India, and Prof. Dr. Mohd. Zulkifli Abdullah from Universiti Sains Malaysia, Malaysia, and Prof. Yasuyuku Nemoto, Ph.D. from Ashikaga University, Japan.

Our special thank also goes to Tarzan Photo and PT. Astaguna Wisesa as our patrons. Also to all individuals and organizations such as the members of international editorial board, the conference organizers, the reviewers and the authors, for their contribution in making TICATE 2019 as a successful international conference and a memorable gathering event. I am also grateful for the support of publication service of IOP Publisher. We hope that the conference could present you wonderful memories to bring home in addition to new insights and friendship congregated during the event.

We truly value your participation and support for the conference. We hope that you will enjoy TICATE 2019 and culture and tradition in Jakarta.

Dr. Hugeng, S.T., M.T. (SMIEEE)



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

**2nd TICATE 2019**  
**Conference Organisation**

**INITIATOR & ORGANIZING INSTITUTION**  
Universitas Tarumanagara, Jakarta

**Supporting Institution**  
IOP Publisher, Tarzan Photo, PT. Astaguna Wisesa

**Honorary Chair**

Prof. Dr. Agustinus Purna Irawan  
R. M. Gatot Soemartono, Ph.D.

**Chairman**

Dr. Hugeng, S.T., M.T. (SMIEEE)  
**Co-chairperson**

Dr. Fransisca Iriani Roesmala Dewi

**Secretary**

Bagus Mulyawan, M.M.

**Program & Sponsorship**

Mei Ie, M.M.

Herlina Budiono, M.M.

**Parallel & Scientific Session**

Dr. Hetty Karunia Tunjungsari  
Sinta Paramita, M.A.

**Treasurer**

Wulan Purnama Sari, M.Si.

**Keynote Speakers**

Prof. Dr. Tjokorda Gde Tirta Nindhia, Universitas Udayana, Indonesia

Dr.-Ing. Joewono Prasetyo, Universiti Tun Hussein Onn, Malaysia

Prof. Dr. Srikantappa A.S., Cauvery Institute of Technology, India

Prof. Dr. Mohd. Zulkifli Abdullah, Universiti Sains Malaysia, Malaysia

Prof. Yasuyuki Nemoto, Ph.D., Ashikaga University, Japan

**Editorial Board / Reviewers:**

Prof. Dr. rer. nat. Alexander Ferrein, University of Applied Sciences Aachen, Germany

Dr.-Ing. A. Ruggeri Toni Liang, Karlsruhe Institute of Technology, Germany

Dr. -Ing Stephan Herzog, TU Kaiserslautern, Germany

Dr. Thomas Marconi, Inside Secure, France

Prof. Yifan Chen, Ph.D., University of Waikato, New Zealand

Dr. Soh Sie Teng, Curtin University, Australia

Dr. Channing Chuang, Kun Shan University, Taiwan

Prof. Mohd Zulkifli bin Abdullah, Universiti Sains Malaysia, Malaysia

Prof. Zaidi Mohd. Ripin, Universiti Sains Malaysia, Malaysia

Dr. -Ing. Joewono Prasetyo, Universiti Tun Hussein Onn, Malaysia

Dr. Filbert H. Juwono, Curtin University, Sarawak Malaysia

Prof. Dr. Tresna P. Soemardi, Universitas Indonesia, Indonesia

Dr. -Ing. Eko Adhi Setiawan, Universitas Indonesia, Indonesia

Prof. Dr. Jamaris, Universitas Gadjah Mada, Indonesia

Dr. Bambang Kismono Hadi, Bandung Institute of Technology, Indonesia

Prof. Eko Sediyyono, Universitas Kristen Satya Wacana, Indonesia

Prof. Tjokorda Gde Tirta Nindhia, Universitas Udayana, Indonesia

Dr. Rianti Ariobimo, Universitas Trisakti, Indonesia

Dr. Richard Napitupulu, Universitas HKBP Nommensen, Indonesia

Prof. Dr. Dyah Erny Herwindiati, Universitas Tarumanagara, Indonesia

Prof. Dr. Leksmono Suryo Putranto, Universitas Tarumanagara, Indonesia

Harto Tanujaya, Ph.D., Universitas Tarumanagara, Indonesia

Jap Tji Beng, Ph.D., Universitas Tarumanagara, Indonesia

Lina, Ph.D., Universitas Tarumanagara, Indonesia

Dr. Steven Darmawan, Universitas Tarumanagara, Indonesia





PAPER • OPEN ACCESS

## Land mapping with least median of squares regression using landsat imagery: a case study Jakarta and sorrounding area

To cite this article: Bagus Mulyawan *et al* 2020 *IOP Conf. Ser.: Mater. Sci. Eng.* **852** 012024

View the [article online](#) for updates and enhancements.

# Land mapping with least median of squares regression using landsat imagery: a case study Jakarta and surrounding area

**Bagus Mulyawan\*, Nicholas Jovianto, Janson Hendryli, and Dyah Erny Herwindati**

Faculty of Information Technology, Universitas Tarumanagara, Jakarta 11440, Indonesia

\*bagus@untar.ac.id; \*Corresponding author's email

**Abstract.** Impervious surface is an area that has undergone natural or semi-natural substitution of artificial land cover which is usually waterproof and relatively permanent, including settlements, industrial buildings, road networks, railroad networks, high voltage electricity networks, airports air and sea ports. Changes cause the narrowing of rivers and open land which become water catchment areas. One of the problems that triggered an increase in impervious land is urbanization. This study aims to map land changes in the Jakarta and surrounding areas using the least median of squares regression method. Based on data from land classification results using 4 classes, Jakarta and Tangerang are areas that have more impervious land than non-impervious land. Meanwhile, Bogor, Bekasi, and Depok are areas that have more non-impervious land.

## 1. Introduction

Impervious surface (built up land) is an area that has undergone natural or semi-natural land cover substitution which is artificial cover which is usually waterproof and relatively permanent, including settlements, industrial buildings, road networks, railroad networks, high voltage electricity networks, airports air and sea ports [2]. According to Stankowski (1972) and Arnold & Gibbons (1996), population growth and urbanization are closely related to an increase in the percentage of developed land that cannot absorb water. Expansion of residential areas can result in a decrease in open land to build land that cannot absorb water. Flood disaster in urban areas is one of the real examples of the decline in open land as a water catchment area. In addition to flooding, various other environmental problems can arise as a result of the change in open land, including erosion disasters, landslides and water crises.

## 2. Method

The existence of outliers in linear regression which is a problem because outliers can cause the formation of a regression parameter model to be less accurate. In overcoming this problem, statisticians try to find alternative estimates of other parameters that are better at overcoming outliers. One of the methods suggested in muscular regression is Least Median of Squares (LMS). Least Median Squares (LMS) is a parameter estimation method with a high breakdown point that was introduced by Peter J. Rousseeuw in 1984. This method predicts the regression parameters by minimizing the median square of the remainder of the observations. The LMS estimator is defined as follows:

$$\hat{b}_{LMS} = \arg \min Q_{LMS}(b). \quad (1)$$

where  $Q_{LMS}(b)$  is the median squared remaining of  $h$  observations,  $\text{med}(e_h^2)$ .



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

In general, the algorithm for applying the LSM Regression method can be summarized in the steps below. Suppose that a data group of size  $N$  is given, and you want to expect a vector  $\theta$  of dimension  $p$  that contains the parameters of the data group. The steps that need to be done are:

1. Determine the size of the subset  $n$ , the number of subsets  $M$  according to the number of classes.
2. Randomly, take  $M$  subset size  $n$  from sample size  $N$ . Find the estimated parameter  $\hat{\theta}_j$  for each subset. Find the median squared error  $e_{ij}^2$  from each subset. Index  $i$  is the index for the example, and index  $j$  is the index for the subset.
3. Define:

$$m = \arg \min [med(e_{ij}^2)] \quad (2)$$

So the subset  $\hat{\theta}_m$  is a subset with the median least square error and  $e_{im}$  is the error vector generated by that subset.

4. Count:

$$S_0 = 1.4826 \left(1 + \frac{5}{N-p}\right) \sqrt{med_i e_{im}^2} \quad (3)$$

5. Calculate the weight of  $w_i$ , for example with  $w_i = 1$ ,  $\left| \frac{e_i}{S_0} \right| \leq median(e)$  and  $w_i = \frac{S_0}{|e_i|}$ .

6. Give  $w_i$  the weight of each example.

7. Make a fit using the weighted least squares method using  $wi$  as the weight to get the final  $\hat{\theta}$ .

The method used in building this application is the least median of square regression method. This method is a regression method that is very suitable to be applied to a calculation of the regression equation model when the data held have outliers. This method will form a regression parameter based on the median value of the data so that outliers will not be very influential on calculations. making this application program will require input data in the form of spatial data in the form of Landsat 8 satellite imagery for the Greater Jakarta area. Landsat 8 satellite imagery data that will be used on the system can be downloaded on the USGS Global Visualization Viewer website page (<https://glovis.usgs.gov/>). The Landsat 8 imagery downloaded will include two intersection files (map codes 122064 and 122065) for each time to cover the whole Jabodetabek area[3]. Jabodetabek map will be used as a subject in the making of an application program to detect changes in land use. The map that is used as data in making this land use change detection application program will cover band 1, band 2, band 3, band 4, and band 5. The choice of using these bands is based on the spectrum represented by each of these bands. Band 1 represents the coastal / aerosol spectrum, making it suitable for detecting territorial waters. Band 2, band 3, and band 4 represent RGB images that are easily captured by the human eye. Band 5 representing the Near-Infrared (NIR) spectrum is commonly used to detect vegetation density when combined with band 4, for example by implementing it in NDVI or SAVI calculations.

### 3. Result and Discussion

Testing with 4 (impervious land, green land, water, and vacant land) class will use testing data for the Greater Jakarta area in the 2014, 2015 and 2017 periods. Comparison between the actual area and the number of pixels per Jabodetabek scene in Landsat 8. can be seen in Table 1.

**Table 1.** Number of Pixels Represented for Jakarta and Surrounding Area

City	Area (km <sup>2</sup> )	Number of pixels per scene
Jakarta	659.26	732,514
Bogor	3,102.35	3,447,053
Depok	180.01	200,015
Tangerang	1,353.45	1,503,838
Bekasi	1,483.90	1,648,775

In image detection using Landsat 8 data, it is known that the resolution of each pixel in the Landsat 8 data represents the original area with a size of 30x30 meters. So that every 1 pixel of Landsat 8 data will represent an actual area of 0.0009 km<sup>2</sup>. The results of the calculation of land area based on each region in 2014, 2015, and 2017 with observations of 4 classes can be seen in Table 2.

**Table 2.** Calculation of Land Area for 4 Classes in 2017

City	Impervious (km <sup>2</sup> )	Green (km <sup>2</sup> )	Water (km <sup>2</sup> )	Clear (km <sup>2</sup> )	Area (km <sup>2</sup> )
Jakarta	570.2382	76.6107	11,7153	0.6984	659.26
Bogor	302,1255	2,460.9663	17,7795	321.4764	3,102.35
Depok	82.1691	95.0427	0,9243	1.8774	180.01
Tangerang	706.6944	581.7204	50,1795	14.8599	1,353.45
Bekasi	697.8627	596.8539	53,2593	135.9216	1,483.90
Jabodetabek	2,359.0899	3,811.1940	133,8579	474.8337	6,778.98

If categorized as impervious and non-impervious land (green land, water, and vacant land), the comparison of the percentage of impervious and non-impervious land detection results for each region based on 4 class observations can be seen in Table 3.

**Table 3.** Percentage of Class Impervious and Non-Impervious 4 Land Detection Results

Years	City	Impervious	Non-impervious
2014	Jakarta	84.78 %	15.22 %
2015	Jakarta	77.54 %	22.46 %
2017	Jakarta	86.49 %	13.51 %
2014	Bogor	8.06 %	91.94 %
2015	Bogor	10.26 %	89.74 %
2017	Bogor	9.75 %	90.25 %
2014	Depok	43.84 %	56.16 %
2015	Depok	39.87 %	60.13 %
2017	Depok	45.64 %	54.36 %
2014	Tangerang	52.41 %	47.59 %
2015	Tangerang	58.11 %	41.89 %
2017	Tangerang	52.21 %	47.79 %
2014	Bekasi	36.23 %	63.77 %
2015	Bekasi	49.45 %	50.55 %
2017	Bekasi	47.02 %	52.98 %

Based on data from land classification results using 4 classes, Jakarta and Tangerang are areas that have more impervious land than non-impervious land. Meanwhile, Bogor, Bekasi, and Depok are areas that have more non-impervious land. Percentage of the amount of impervious and non-impervious land every year experiences fluctuating changes. In the Bogor, Tangerang and Bekasi regions, the number of impervious land has increased in 2015 and decreased in 2017. While in the Jakarta and Depok area, impervious land has decreased in 2015 and experienced an increase in 2017.

#### 4. Conclusion

In general, the most frequent changes in land use in most Jabodetabek areas are land use changes from the type of green land to impervious land. The following are some regional data samples that experienced changes in land use in the period 2014 to 2017 which can be seen from Figure 1 and Figure 2.




Figure 1. Serpong Area (2014 and 2017)




Figure 2. Tangerang Area (2014 and 2017)

## 5. References

- [1] Herwindiati, Dyah E., Hendryli, Janson, & Hiryanto, Lely. (2017). Impervious Surface Mapping Using Robust Depth Minimum Vector Variance Regression. *European Journal of Sustainable Development* **6(3)** 29-39
- [2] Rousseeuw, P. J. (1984). Least Median of Squares Regression. *American Statistical Association*, 79(388), 871-880
- [3] U. S. Geological Survey. (2016). Landsat 8 Data User Handbook. Retrieved from <https://landsat.usgs.gov/sites/default/files/documents/Landsat8DataUsersHandbook.pdf>
- [4] Draper, N. R. & Smith, Harry. (2014). *Applied Regression Analysis, Third Edition*. New York: John Wiley
- [5] Rousseeuw, P. J. & Leroy, A.M. (1987). *Robust Regression and Outlier Detection*. New York: John Wiley