

eISSN 2798-1630

EBERS PAPYRUS

Jurnal Kedokteran & Kesehatan

VOL. 27 No. 1 JUNI 2021



GEDUNG
DR. H.R. SOEWONDO

Fakultas Kedokteran Universitas Tarumanagara

DEWAN REDAKSI

Ketua Editor	Ria Buana	
Editor	Erick Sidarta Fia Fia Lie Hari Darmawan Octavia Dwi Wahyuni	
Mitra Bebestari	Herwanto Meilani Kumala Oentarini Tjandra Shirly Gunawan Triyana Sari Twidy Tarcisia Wiyarni Pambudi	(Universitas Tarumanagara) (Universitas Tarumanagara) (Universitas Tarumanagara) (Universitas Tarumanagara) (Universitas Tarumanagara) (Universitas Tarumanagara) (Universitas Tarumanagara)
Sekretariat	Clement Drew	
Alamat Redaksi	Fakultas Kedokteran Universitas Tarumanagara Jl. Letjen S.Parman No. 1 Jakarta Barat 11440 Telp: 0215671781 <i>email:</i> jurnal.eberspapyrus@gmail.com	

EBERS PAPYRUS

Jurnal Kedokteran & Kesehatan

Vol. 27 No. 1 Juni 2021

DAFTAR ISI

KULIT MUKA			
1	Henrik Blum	Andri Wanananda	1-3
TINJAUAN PUSTAKA			
2	Peranan Bahan Alam Dalam Pandemi Covid-19 The Role Of Natural Materials In The Covid-19 Pandemic	Fenny Yunita	4-15
ARTIKEL ASLI			
3	Comparison Of Breastmilk Production From Mothers Of Premature And Mature Babies During The First Week In Nicu Siloam Hospitals Kebon Jeruk	Naomi Esthernita F. Dewanto, Nancy Napitupulu	16-25
4	Hubungan Antara Pengetahuan Akseptor Kb Suntik Dengan Ketepatan Waktu Suntik Kb Di Bps Khanifah, Kabupaten Tangerang, Propinsi Banten Periode Januari – Desember 2017	Riffany Krisdiana, Ernawati	26-33
5	Determinan Persepsi Dan Perilaku Ibu Dalam Membaca Label Makanan Ringan Anak Tk-Sd Bumi Serpong Damai	Ruth Brigitta Salim, Wiyarni Pambudi	34-44
6	Gambaran Kadar Glukosa Darah Pada Lansia Di Sentra Vaksinasi Covid-19 Fakultas Kedokteran Universitas Tarumanagara Jakarta	Dorna Yanti Lola Silaban	45-53
7	Gambaran Jumlah Trombosit Pada Mahasiswa Fakultas Kedokteran Universitas Tarumanagara Angkatan 2012-2016 Dengan Obesitas	Daniel Albar R Patterson, Sari Mariyati Dewi	54-61
8	Faktor-Faktor Yang Mempengaruhi Pemberian Susu Formula Pada Bayi Usia 6 Bulan Di Posyandu Bougenville Jakarta Barat Tahun 2020	Nathaza Caroline Ipsan, Naomi Esthernita Fauzia Dewanto\	62-74
9	Korelasi Faktor Demografi, Nutrisi Dan Antropometri Terhadap Kejadian Hipertensi Pada Petugas Keamanan Di Universitas Tarumanagara Jakarta	Alexander Halim Santoso, Susy Olivia Lontoh, Farell Christian Gunaidi, Ferdinand Harum	75-90

		Istikanto, Edwin Destra, Junius Kurniawan, Natasha Anggraeni, Stanley Surya Cahyadi, Tizander Mayvians	
10	Pemetaan Faktor-Faktor Risiko Sindroma Metabolik Pada Mahasiswa Fakultas Kedokteran Universitas Tarumanagara Jakarta	Victor Sepriyadi Wohangara, Alexander Halim Santoso	91-99
11	Penelitian Pendahuluan Tingkat Kebugaran Fisik Dengan Six Minute Walk Test Pada Mahasiswa Fakultas Kedokteran Universitas Tarumanagara	Irene Fransisca Gunalam, Susy Olivia Lontoh	100-115

LAPORAN KASUS

12	Perdarahan Iliopsoas Pada Pasien Covid-19 Dalam Terapi Antikoagulan	Andria Priyana	116-123
----	---	----------------	---------

COMPARISON OF BREASTMILK PRODUCTION FROM MOTHERS OF PREMATURE AND MATURE BABIES DURING THE FIRST WEEK IN NICU SILOAM HOSPITALS KEBON JERUK

Oleh:

Naomi Esthernita F. Dewanto¹, Nancy Napitupulu²

¹Pediatric Department, Medical Faculty, Tarumanagara University, Jakarta

2 Siloam Hospital, Jakarta

Correpondence: naomiesthernita@gmail.com

Abstract

Background: The principal goal for infants especially preterm is the provision of the mother's own milk (MOM), but on the other hand it is difficult to get MOM for preterm babies due to delayed of lactogenesis II and mother's stress. Milk production and adequacy, for mothers of both healthy breastfeeding term infants and non-nursing preterm infants, have been shown to have a significant relationship with milk production 4-6 days after birth. This research aim to describe the comparison of milk production between mothers of premature and mature babies during the first week in NICU SHKJ.

Methods: This is an observational descriptive with comparative study of mother's milk production of preterm (<37 weeks) and term (≥ 37 weeks) babies during the first week of life in NICU Siloam Hospitals Kebon Jeruk from 1st January until 31st December 2018. We documented the milk production from mothers breastmilk expression every 3-4 hours since 4-6 hours after delivery until day 7.

Results: From 181 babies during 2018, we got 31 preterm and 77 term babies that we could documented from birth until day 7. The average breastmilk production of the preterm vs term mothers in each expression from day 1-7 were 0,81 vs 1,61 ml; 3 vs 3,65 ml; 6,66 vs 6,94 ml; 21,5 vs 26,6 ml; 34,1 vs 46,4 ml; 45,92 vs 46 ml; 48,88 vs 50,4 ml/x expression.

Conclusions: Expressing milk for a preterm or unwell baby requires commitment by the mother and effective support from staff. Commence breast milk expression as soon as possible after birth, at least within the first six hours, followed by frequent, regular and effective breast milk expression to stimulate adequate breast milk production may provide MOM for babies even for preterm babies in NICU with almost similar amount.

Keywords: Breastfeeding, mothers own milk, milk expression, NICU

Introduction

Human milk is considered the best nutrition for new-born infants because it provides substantial nutritional, cognitive, emotional, and immunologic benefits for the infant.^{1,2} Breastfeeding confers protection against gastrointestinal infections, respiratory infections, otitis media, allergic diseases, reduces risk of Sudden Infant Death Syndrome (SIDS) and it is also associated with a reduced long term risk of diseases such as inflammatory bowel disease (IBD), celiac disease, obesity, diabetes, leukemia and lymphoma.³⁻⁸ The protective role of human milk seems to be the consequence of a synergistic action of the wide range of health-promoting components such as carbohydrates, nucleotides, fatty acids, immunoglobulins, cytokines, immune cells, lysozyme, lactoferrin and other immunomodulatory factors.⁹⁻¹² All of the benefits of human milk are magnified in the preterm infant as it protects against a multitude of NICU challenges.^{2,13} Early breast milk, known as colostrum, provides immune benefits to infants, offering potential risk reduction for nosocomial infection, sepsis and necrotizing enterocolitis (NEC).¹⁴ It lowers the mortality rates, decrease length of hospital stay and long-term growth failure and improves feeding tolerance. Other benefits include a decreased readmissions after NICU discharge and maternal psychological benefits.¹⁵⁻²⁰ Long- term studies also suggest that human milk feeding in NICU is associated with greater neurodevelopmental outcomes, lower rates of severe Retinopathy of Prematurity (ROP), metabolic syndrome, blood pressures and LDL concentrations and improved leptin and insulin metabolism.^{8,21,22}

Several studies indicate that immunoprotective factors are more highly concentrated in the colostrum of mothers who deliver preterm infants than those who give birth at term.^{23,24} Provision of colostrum is recognized as oral immune therapy (OIT) and is valuable to all NICU infants unable to feed orally including extremely low birth weight (ELBW; BW<1000g) infants.^{25,26} During this intervention, a small amount of colostrum is placed directly onto the oropharyngeal mucosa for absorption.²⁷ In theory, the abundant immune factors in colostrum interact with lymphoid tissues in the oropharynx and stimulate the immature neonatal immune system.²⁸

Mothers of preterm infants experience physiological and emotional challenges, which may adversely affect breast feeding outcome.²⁹ The reasons are:

- mammary growth may be incomplete;
- decreased exposure of mammary epithelium to prolactin, cortisol, and other hormones that normally occur during a full-term pregnancy ;
- poor placental function with low levels of placental lactogen.³⁰

Maternal stress and several causes underlying preterm birth (eg, hypertension, diabetes, and maternal obesity) have been associated with delayed lactogenesis II, which begins with the production of copious amounts of breastmilk following delivery.^{20,30-33} These mothers start and sustain breast feeding at lower rates than mothers of term babies.³⁰ A delay in lactogenesis stage II also common in term infants, especially if initiation of breastfeeding is delayed following delivery.³⁴ However,

some mothers seem to achieve compensatory growth by the adoption of a frequent milk expression schedule in the early postpartum period.^{29,35,36} The first milk expression should take place as soon as possible (within 1 hour) following delivery.³⁶⁻³⁸ The amount produced can be as little as a few drops of colostrum at each expression for the first 24–48 hours post partum. Mothers should be encouraged to express at least eight times in 24 hours.³⁹ Milk production and adequacy (defined as greater than 500 ml/day), for mothers of both healthy breastfeeding term infants and non-nursing preterm infants, have been shown to have a significant relationship with milk production 4–6 days after birth.^{28,40} Because of the abundance benefit of breastmilk for babies especially in NICU, we want to report the use of MOM and compare the breastmilk production from mothers of preterm and term babies in our NICU.

Material & Methods

This is an observational descriptive with comparative study of mother's milk production of preterm (<37 weeks) and term (≥ 37 weeks) babies during the first week of life in NICU Siloam Hospitals Kebon Jeruk from 1st January until 31st December 2018. We have admitted 181 patients in NICU during 2018 consist of 43 preterm and 138 term babies, but we can only documented 31 samples from preterm and 77 samples from term babies. The uncomplete breastfeeding documentation was due to not all the babies were born in our NICU, so they were referred from other hospital not at the first day of life. Another reasons were mom didn't express breastmilk regularly even though we already encourage them to do so or administration mistakes by NICU staff.




Figure 1. Sampling design

We encourage all mothers in our NICU to express MOM. If they were give birth in our hospital, our policies is to express breastmilk after 4-6 hours of delivery. We teach the parents to wash hands before expressing breastmilk and how to express breastmilk with hands (using syringe) or breast pump. Then, we try to get breastmilk expression every 3-4 hours, collect the data for 1 week (from colostrum until day 7 everyday) and fill the amount of breastmilk in the form below:

Lembar Pemantauan Produksi Air Susu Ibu (ASI)

Identitas Pasien :							
Diagnosis :							
Usia Gestasi :							
Pompa (Jam)	Jumlah ASI (dalam satuan ml)						
	Hari I	Hari II	Hari III	Hari IV	Hari V	Hari VI	Hari VII
TOTAL							

Figure 2. Breastmilk production documentation form

Results

The characteristic of our samples are shown in the table below:

Table 1. Sample Characteristics

	Preterm Babies		Term Babies	
	n=3 1	%	n=7 7	%
Gestational Age				
<32 week	4	12,9		
≥32 -35 week	15	48,4		
≥35 -37 week	12	38,7		
≥37 week			77	100

Number of Pregnancies				
G1	19	61,3	37	48
G2	8	25,8	24	31,2
G3	4	12,9	12	15,6
≥G4	0	0	4	5,2

Delivery Method				
Vaginal Delivery	4	12,9	16	20,8
Vacuum Extraction	0	0	1	1,3
Caesarean Section	27	87,1	60	77,9

Baby Gender				
Boy	18	58,1	47	61
Girl	13	41,9	30	39

Birth Weight				
1000 - <1500 g	5	16,1	0	0
1500 - <2500 g	21	67,8	6	7,8
2500 - <4000 g	5	16,1	67	87
≥ 4000 g			4	5,2

We calculate the average amount of breastmilk produced by each mom everyday then we differentiate the average from all preterm compare to term babies from day 1 until day 7 of birth. The chart is presented in figure 3.




Figure 3. Breastmilk Production Chart

The breastmilk production is increasing everyday both in preterm and term babies. The highest increment can be seen in day 4 after birth and continues until day 7. Breastmilk production of preterm babies seem to be fewer than term babies in day 1 until day 5, but become almost same in day 6 until day 7 after birth. The highest average breastmilk production per expression in our NICU is 50.4 ml for term baby and 48.8 ml for preterm baby at day 7.

Discussion

Human milk is the preferred food for infants, including ill and preterm infants. As we all know, it is difficult to get MOM for preterm babies due to delayed of lactogenesis II and mother's stress. Because of that, we try to implement the 5 concepts of Family Centred Care (FCC) to support the successful of MOM production in our NICU. It include family's care taking, equal family participation, collaboration, maintaining family's respect and dignity & knowledge transformation.⁴¹

The implementation of FCC concepts is as follows:

1. Change in paradigm of care-giving in NICU
2. Parents are integral part of the care team and are partners with nurses and medical team
3. Nurses are teachers and consultants to parents
4. Parents assume most of the care of baby, except for IV, medication and tests
5. Parents gain confidence, knowledge and control.⁴¹

We build the FCC concept and special rooms in our NICU so mothers can stay 24 hours beside the babies. They can do Kangaroo Mother Care (KMC) and breastfeed with Cpap. Mothers are encouraged to empty the breast every 3 hours using hands for 3 initial days and breastpump thereafter. NICU staff and Lactation team routinely educate and monitor the discipline of the mothers. We also give psychological support to reduce maternal stress. We document breastmilk production per expression from day 1 until day 7 after birth. We adjust the lactation intervention according to patient's needs e.g. breast massage, lactation massage, lactagogue or supplementation with breastmilk donor.

Our hospital especially NICU staff and Lactation team always encourage mothers of ill and premature babies to begin milk expression within 4-6 hours following delivery. This mothers produced greater milk volumes at the initial milk expression session and at every 24-hour period during the first 7 days following delivery. Breast milk production between mothers of premature and mature babies in the first 7 days of life in our NICU is not significantly different.

Mothers who began milk expression after 6 hours following delivery had an earlier onset of lactogenesis stage II. Lactogenesis stage II is defined by a sudden feeling of breast fullness. In term mothers, timing of lactogenesis stage II begins by approximately 36 hours following delivery and is completed by approximately 4 days. However, 82% of mothers delivering premature infants had an impaired lactogenesis stage II and there was a finding reported a 1-day delay in the onset of lactogenesis stage II in mothers of premature infants.²⁰ In this study, mothers experienced a delay in timing of lactogenesis stage II, which begins at day 4 after birth.

Hill et al. found an association between timing of expression initiation and adequate milk production (defined as milk output of ≥ 3.500 mL/week) in weeks 2-6.²⁰ In this study, the highest average breastmilk production per expression in our NICU is 50.4 ml, around 2822.4 mL/week in week 1. This data support 63.2% successful rate of exclusive breastfeeding in our NICU during 2018.

Conclusions

Expressing for a preterm or unwell baby requires commitment by the mother and effective support from staff. Commence breast milk expression as soon as possible after birth and at least within the first six hours, followed by frequent, regular and effective breast milk expression to stimulate adequate breast milk production may provide MOM even for preterm babies in NICU.

References

1. Anderson J. W., Johnstone B. M., Remley D. T. Breast-feeding and cognitive development: a meta-analysis. *Am J ClinNutr* 1999;70:525–35.
2. Lawrence R. M., Lawrence R. A. Breastfeeding: More Than Just Good Nutrition. *Pediatrics in Review* 2011;32:267-279.
3. DuijtsL., Jaddoe V.W., Hofman A., Moll H.A. Prolonged and exclusive breastfeeding reduces the risk of infectious diseases in infancy. *Pediatrics* 2010;126: e18-25.
4. Ip S., Chung M., Raman G., Chew P., Magula N., DeVine D., Trikalinos T., Lau J. Breastfeeding and maternal and infant health outcomes in developed countries. *Evidence Report/ Technology Assessesment* 2007;153: 1-186.
5. Greer F.R., Sicherer S.H., Burks A.W. Effects of early nutritional interventions on the development of atopic disease in infants and children: the role of maternal dietary restriction, breastfeeding, timing of introduction of complementary foods, and hydrolyzed formulas. *Pediatrics* 2008; 121: 183-191.
6. Hauck F. R., Thompson J. M. D., Tanabe K. O., Moon R. Y., Vennemann M. M. Breastfeeding and Reduced Risk of Sudden Infant Death Syndrome: A Meta-analysis. *Pediatrics* 2011; 128(1): 1-8.
7. Stolzer J.M. Breastfeeding and obesity: a meta-analysis. *Open J Prev Med* 2011; 1(3): 88-93.
8. American Academy of Pediatrics. Breastfeeding and the Use of Human Milk. *Pediatrics* 2012; 129(3): e827-41.
9. Van 't Land B., Boehm G., Garssen J. Breast milk: components with immune modulating potential and their possible role in immune mediated disease resistance. In: Watson R.R., Zibadi S., PreedyV.R. (eds.) *Dietary components and immune function*. Springer, Berlin, Germany. 2010;pp. 25-41.
10. Jeurink P. V., Bergenhennegouwen J., Jimenez E., Knippels L. M. J., Fernandez L., Garssen J., Knol J., Rodriguez J. M., Martin R. Human milk: a source of more life than we imagine. *Beneficial Microbes*, March 2013; 4(1): 17-30.
11. Cacho N. T., Lawrence R. M. Innate Immunity and Breast Milk. *Front. Immunol* 2017;8:584.
12. Fernandez L., Langa S., Martin V., Maldonado A., Jimenez E., Martin R., Rodriguez J. M. The human milk microbiota: Origin and potential roles in health and disease. *Pharmacological Research* 2013;69:1–10
13. Maffei D., Schanler R. J. Human milk is the feeding strategy to prevent necrotizing enterocolitis! *Seminars in Perinatology* 2016: 1-5.
14. Patel A. L., Johnson T. J., Engstrom J. L., et al; Impact of Early Human Milk on Sepsis and Health Care Costs in Very Low Birth Weight Infants. *J Perinatol* 2013; 33(7): 514–519.
15. Vohr B. R., Poindexter B. B., Dusick A.M. et al. Beneficial Effects of Breast Milk in the Neonatal Intensive Care Unit on the Developmental Outcome of Extremely Low Birth Weight Infants at 18 Months of Age. *Pediatrics* 2006;118(1): e115-23.

16. Vohr B. R., Poindexter B. B., Dusick A. M., et al; National Institute of Child Health and Human Development National Research Network. Persistent beneficial effects of breast milk ingested in the neonatal intensive care unit on outcomes of extremely low birth weight infants at 30 months of age. *Pediatrics* 2007;120(4):e953-9.
17. Schanler R. J., Shulman R. J., Lau C. Feeding strategies for premature infants: beneficial outcomes of feeding fortified human milk versus preterm formula. *Pediatrics* 1999;103(6 pt 1):1150–1157.
18. Hintz S. R., Kendrick D. E., Stoll B. J., et al; NICHD Neonatal Research Network. Neurodevelopmental and growth outcomes of extremely low birth weight infants after necrotizing enterocolitis. *Pediatrics* 2005;115(3):696–703.
19. Sweet L. Expressed breast milk as 'connection' and its influence on the construction of 'motherhood' for mothers of preterm infants: A qualitative study. *Int Breastfeed J* 2008;3:30–42.
20. Parker L. A., Sullivan S., Krueger C., Mueller M. Association of Timing of Initiation of Breastmilk Expression on Milk Volume and Timing of Lactogenesis Stage II Among Mothers of Very Low-Birth –Weight Infants. *Breastfeeding Medicine* 2015;10(2):84–91.
21. Isaacs E. B., Fischl B. R., Quinn B. T., Chong W. K., Gadian D. G., Lucas A. Impact of breast milk on intelligence quotient, brain size, and white matter development. *Pediatr Res* 2010;67(4):357-362
22. Lucas A. Long-term programming effects of early nutrition—implications for the preterm infant. *J Perinatol* 2005;25(suppl 2):S2–S6.
23. Lee J., Kim H. S., Jung Y. H., et al; Oropharyngeal Colostrum Administration in Extremely Premature Infants: An RCT. *Pediatrics* 2015;135(2): e357-66.
24. Dvorak B., Fituch C. C., Williams C. S., Hurst N. M., Schanler R.J. Increased epidermal growth factor levels in human milk of mothers with extremely premature infants. *Pediatr Res* 2003;54(1):15–19.
25. Pletsch D., Ulrich C., Angelini M., Fernandes G., Lee D. S. C. Mother' "Liquid Gold": A Quality Improvement Initiative to Support Early Colostrum Delivery via Oral Immune Therapy (OIT) to Premature and Critically Ill Newborns. In: A Quality Improvement Initiative to Support Early Colostrum Delivery. *Nurs. Leadersh (Tor Ont)* 2013;26: 34-42.
26. Rodriguez N. A., Groer M. W., Zeller J. M., et al; A Randomized Controlled Trial of the Oropharyngeal Administration of Mother's Colostrum to Extremely Low Birth Weight Infants in the First Days of Life. *Neonatal Intensive Care* 2011;24(4):31-35.
27. Rodriguez N. A., Meier P. P., Groer M. W., Zeller J. M. Oropharyngeal administration of colostrum to extremely low birth weight infants: theoretical perspectives. *J Perinatol* 2009;29(1):1–7.
28. Gephart S. M., Weller M. Colostrum as oral immune therapy to promote neonatal health. *Adv Neonatal Care* 2014;14(1):44–51.
29. Jones E., Spencer S. A. OptimisingThe Provision of Human Milk For Preterm Infants. *Arch Dis Child Fetal Neonatal Ed.* 2007 Jul;92(4):F236-F238.
30. Dewey K.G. Maternal and Fetal Stress Are Associated with Impaired Lactogenesis

in. *Humans. J. Nutr* 2001;131:3012S-3015S.

31. Wilson E., Christensson K. Early Provision of Mother's Own Milk and Other Predictors of Successful Breast Milk Feeding after Very Preterm Birth: A Regional Observational Study. *J Hum Lact* 2015;1-8.
32. Matias S. L., Dewey K. G., Quesenberry C. P. Jr., Gunderson E. P. Maternal prepregnancy obesity and insulin treatment during pregnancy are independently associated with delayed lactogenesis in women with recent gestational diabetes mellitus. *Am J ClinNutr* 2014;99(1):115-121.
33. Cregan M. D., De Mello T. R., Kershaw D., McDougall K., Hartmann P. E. Initiation of lactation in women after preterm delivery. *Acta Obstet Gynecol Scand* 2002;81(9):870- 877.
34. Dewey K. G., Nommsen-Rivers L. A., Heinig M.J., et al. Risk factors for suboptimal infant breastfeeding behavior, delayed onset of lactation, and excess neonatal weight loss. *Pediatrics* 2003;112:607-619.
35. Parker L. A., Sullivan S., Krueger C., Kelechi T., Mueller M. Effect of early breast milk expression on milk volume and timing of lactogenesis stage II among mothers of very low birth weight infants: A pilot study. *Journal of Perinatology* 2012;32:205-209.
36. Spatz D., Edwards T. The Use of Human Milk and Breastfeeding in the Neonatal IntensiveCare Unit. National Association of Neonatal Nurses. 2015.
37. Schanler R. J., Hurst N. M., Lau C. The use of human milk and breastfeeding in preterm infants. *ClinPerinatol* 1999;26:379-98.
38. Furman L., Minich N., Hack M. Correlates of lactation in mothers of very low birthweight infants. *Pediatr* 2002;109:e57.
39. Neville M. C., Morton J., Umemura S. Lactogenesis, the transition from pregnancy to lactation. *PediatrClin North Am* 2001;48:35-52.
40. Kent J.C., Gardner H. Breastmilk Production in the First 4 Weeks after Birth of Term Infants. *Nutrients* 2016;8(12):756.
41. Ramezani T., Shirazi Z. H., Sarvestani R. S., Moattari M. Family-Centered Care in Neonatal Intensive Care Unit: A Concept Analysis. *IJCBNM* 2014;2:4.