

Basil_Leaf_Extract_Ocimum_x_afri canum_Lour

by Limanan David

Submission date: 09-Jan-2024 03:27PM (UTC+0700)

Submission ID: 2268296655

File name: f_Basil_Leaf_Extract_Ocimum_x_afri
canum_Lour_-_david_limanan.pdf (416.24K)

Word count: 3126

Character count: 16775

32

Phytochemical Screening, Total Antioxidant Capacity, and Toxicity Test of Basil Leaf Extract (*Ocimum x africanum* Lour.).

Timotius^{1*}, David Limanan², Eny Yulianti² and F. Ferdinal²

12

¹ Faculty of Medicine, Tarumanagara University, Jakarta, Indonesia

² Staff of Department of Biochemistry and Molecular Biology, Faculty of Medicine, Tarumanagara University, Jakarta, Indonesia

*Corresponding author. Email: timotius.405180098@stu.untar.ac.id

ABSTRACT

High pollutant levels have caused an increase in disease that correlated with Reactive Oxygen Species through a condition that is oxidative stress. Oxidative stress can be stabilized with antioxidants by stabilizing active and unstable radicals to become inactive and stable. Basil leaves (*Ocimum x africanum* Lour) are known to have antioxidant capacity. The aim of this study is to determine the phytochemical profiling and antioxidant capacity of basil leaves extract. This research was conducted with an in vitro experimental study design. Extraction of this research was carried out by maceration method using methanol as solvent. The tests were carried out in the form of phytochemical tests to determine the content of secondary metabolites and antioxidant capacity tests using the DPPH method. In phytochemical test of basil leaf extract, 16 was found positive for secondary metabolites of alkaloids, anthocyanins, cardio glycosides, coumarins, Flavonoid, glycosides, phenols, quinones, saponins, steroid, terpenoids and tannins and the antioxidant capacity test obtained IC_{50} of 174.04 μ g/mL. In this study we found that basil leaves contain secondary metabolites and antioxidants capacity but not as much as vitamin C, even though basil leave antioxidant capacity not as much as vitamin C, basil leave have other compound like Linalool that have antimicrobial and antioxidant activity, beside vitamin C also can induce acid reflux, so it can be concluded that basil leaves have secondary metabolites and the antioxidant capacity.

13

Keywords: *Ocimum x africanum* Lour, Reactive Oxygen Species, Total Antioxidant Capacity.

1. INTRODUCTION

7

In the past decade, great attention has been paid to the problem of exposure to air pollution due to vehicle traffic and other combustion processes. Air pollutants such as Particulate matter (PM) and gas are considered the most important factors in urban areas [1]. Each air pollutant can cause toxic effects on the respiratory and cardiovascular systems, such as ozone, nitrogen oxides, and suspended particles all of which have the same properties as strong oxidants that are a source of reactive oxygen species (ROS) [2]. Sources of ROS can result from transition metal chemical reactions (iron, copper, and chromium) that catalyze fenton reactions therefore capable of inducing oxidative damage to DNA [3].

Oxidative Stress due to fenton reaction catalyzed by this transition metal can generate hydroxyl radicals that are the source of ROS in the body's system [4], thus disrupting systemic organs such as the lungs, heart, liver and kidneys [5]. As a result the human body system needs defenses against ROS, that is antioxidants [6].

Antioxidants are divided into 2 categories such as *endogenous antioxidants* and *exogenous antioxidants*. *Exogenous antioxidants* can be obtained from nutrients, to support antioxidants that needed by the human body [7]. one of the sources *exogenous antioxidant* derived from Basil (*Ocimum x africanum* lour), methanolic extract content of basil leaves has antioxidant properties that neutralize ROS so that it can be hepatoprotective, Anti-carcinogenesis, and Cardioprotective [8].

3

Copyright © 2021 The Authors. Published by Atlantis Press International B.V.

This is an open access article distributed under the CC BY-NC 4.0 license -<http://creativecommons.org/licenses/by-nc/4.0/>. 52

Basil belongs to the family *Lamiaceae* which is a plant used as a foodstuff. In addition, empirically basil is used for complementary foods, skin diseases, antifungals and antiemetics [9].

Basil leave is a part of basil that known to have a high phenolic content that encourages the author to conduct further research on the antioxidant ability of basil leaves.

2. MATERIAL AND METHOD

This Experimental research using basil leave extract to be test on phytochemical test based on Harborne as reference and antioxidant activity with DPPH method. Basil leave collected as much as 700 grams thereafter this leave be dried to obtain 100 grams of simplicia after that, the simplicia macerated using methanol solvent and then evaporated using rotatory evaporator therefore will produce extract weight as much as 14.59 gram and the yield was 14.59 %. With the equation below:

$$\% \text{Yield} = \frac{\text{Extract weight}}{\text{Dried weight}} \times 100\% \quad (1)$$

$$= \frac{14.59}{100} \times 100\% \quad (2)$$

With the basil leaves extract, we tested the sample to test the phytochemical content that used mayer and dragendroff to find the alkaloid, NaOH to find anthocyanin, betacyanin, and flavonoid, keller-kiliani to find cardio glycosides, NH3 to find coumarin, *Modified Borntrager* to find glycosides, folin ciocalteau to find phenols, H₂SO₄ to find quinone, Foam test to find saponin, Liebermann-Burchard to find steroid, salkowski to find terpenoid, *ferric-chloride* to find tannins.

Antioxidant capacity test (DPPH) using Blois Method. Once the absorbance data of each concentration that has been measured by genesis 30 vis spectrophotometry, the amount of absorption resistance can be calculated through the average absorbance in tubes A and B with the formula:

mean sample absorbance =

$$\frac{\text{Absorbance A} + \text{Absorbance B}}{2} \quad (1)$$

$$2 \quad (2)$$

% Inhibition is a large percentage of the sample's ability to inhibit the absorption of free radicals, which in this case is used as DPPH radicals.

Control absorption is defined as the result of absorption of DPPH solution with a concentration of 50 μM at maximum wavelength. Meanwhile, the average absorbance of the sample is the result of the absorption of the sample in accordance with the concentration of each sample that has been given radical DPPH 50 μM at the maximum wavelength. The graph (*simple*

scatter) is then created according to the data that has been obtained between the concentration (x axis) and %inhibition (Y axis) and note the formation of linear lines, the result of the value R² and its linear regression equation (Y = aX + b). After that data is collected by measuring basil leave extract levels through antioxidant activity of DPPH method. Measurement antioxidant activity used Genesys 30 vis spectrophotometer by Collect the sample absorbance value.

Data are expressed as mean standard error of the mean. a presented in the form of tables and graphs. All statistical analysis w performed using GraphPad Prism v. 9.0 (GraphPad Software, San Diego, CA). p-value of <0.05 was considered statistically significant.

3. RESEARCH RESULTS

Phytochemicals of Basil Leave Extract

Based on the results of qualitative phytochemical tests basil leave extract obtained that this extract contains alkaloids, anthocyanins, cardio glycosides, coumarin, flavonoids, glycosides, phenolic, quinine, saponins, steroids, terpenoids, and tannins (Table 1). But no betacyanin compounds were found.

Total Antioxidant Capacity of Basil Leave

Table 1. Phytochemical Screening of Basil Leaf

Phytochemical	Extract	Method/Reagen
Alkaloid	+	Mayer/Wagner
Antosianin	+	NaOH
Betasianin	-	NaOH
Kardio glikosdia	+	Keller Kiliani
Kumarin	+	NaOH & Kloroform
Flavonoid	+	NaOH
Glikosida	+	Modified Borntrager
Fenolik	+	Foil Ciocalteau
Kuinon	+	H ₂ SO ₄
Saponin	+	Foam
Steroid	+	Liebermann Burchard
Terpenoid	+	Salkowski
Tanin	+	Ferric Chloride

Extract

The absorbance of the test was determined by using ethanol extract *ocimum x africamum lour* weighed as much as 25 mg and dissolved with a solution of methanol in a measuring flask of 25 mL, so that the stock obtained with a concentration of 1000 $\mu\text{g}/\text{mL}$. Then dilute with a solution of methanol so that a concentration of 100, 200, 300, 400 $\mu\text{g}/\text{mL}$ in a measuring flask of 10 mL, then stirred until evenly. To find out the required volume according to the specified concentration, the formula $M_1 \cdot V_1 = M_2 \cdot V_2$ is used. Prepare 4 test tubes and done duplo, so it takes 8 test tubes in total.

Table 2. Antioxidant Capacity of Basil Leaf Extract

Concentration ($\mu\text{g}/\text{mL}$)	Absorbance (516 nm)	Inhibition percentage (%)	IC_{50} ($\mu\text{g}/\text{mL}$)
100	0,431	30,484	174,04
200	0,261	58,548	
300	0,112	88,742	
400	0,071	88,226	

Using aluminum foil to wrap the entire test tube without exception. Prepare a 2 mL microtube tube for each extract sample concentration and then insert it into the tube until it is slightly full for easy sampling when DPPH is added. Into the test tube taken as much as 0.5 mL sample from microtube using micropipettes then inserted into the test tube according to their respective concentrations that have been determined earlier. Then added 3.5 mL of DPPH stock solution, after stirring using vortex until well mixed, then incubated in a dark room with room temperature for 30 minutes. After 30 minutes, the absorbance is searched using a 30 vis genesys spectrophotometer. The calibration curve is based on the relationship between the concentration of basil leaves extract and the results of the calculation of the percentage of inhibition.

Maximum wavelength of DPPH is 516 nm with the maximum absorbance obtained is 0.620 nm. The absorbance value of each basil leave extract concentration (*Ocimum x africamum lour*) is read using a Uv-vis spectrophotometer at a wavelength of 510 nm and is calculated percent (%) inhibition of each concentration from (Table 2) then created curves and searched for linear equations. From the percentage value of inhibition obtained, a straight-line graph will be formed so that there is a linear equation $y = 0.1974X + 15.645$ with $R^2 = 0.933$ (Fig 1). This equation is used to

find IC_{50} capacity of total antioxidant basil leave extract and obtained IC_{50} basil leave extract was 174.04 $\mu\text{g}/\text{mL}$.

Antioxidant Capacity of Vitamin C

The standard antioxidant capacity was determined by using vitamin C (ascorbic acid) by means of ascorbic acid 0.01 grams dissolved in a solution of methanol up to 100 mL. Then a stock with a concentration of 100 $\mu\text{g}/\text{mL}$ will be formed. After that the dilution of ascorbic acid is done by taking stock as much as 0.2 mL, 0.3 mL, 0.4 mL, 0.5 mL, and 0.6 mL in measuring flask and dissolved with methanol up to 10 mL in all five stocks. Obtained concentrations of 2 $\mu\text{g}/\text{mL}$, 3 $\mu\text{g}/\text{mL}$, 4 $\mu\text{g}/\text{mL}$, 5 $\mu\text{g}/\text{mL}$, 6 $\mu\text{g}/\text{mL}$, respectively. Then take 3.5 mL of DPPH and added as much as 0.5 mL of ascorbic acid is piped of each dilution. Performed incubation for 30 minutes in a dark enclosed room at room temperature. After that, the absorbance is read at optimal wavelengths using a 30 vis genesys sp 310 spectrophotometer. The calibration curve is based on the relationship between the concentration of ascorbic acid and the result of the calculation of the percentage of inhibition.

Table 3. Standard Levels of Ascorbic Acid

Concentration ($\mu\text{g}/\text{mL}$)	Absorbance (516 nm)	Inhibition percentage (%)	IC_{50} ($\mu\text{g}/\text{mL}$)
2	0,346	26,85	5,40
4	0,288	39,11	
6	0,213	54,96	
8	0,152	61,86	
10	0,086	81,81	

Each concentration of vitamin C that has been mixed with DPPH is read using uv-vis spectrophotometer (Table 3). Once absorbance is obtained per concentration, it is calculated as percent (%) inhibition then made a standard curve and looked for linear equations. From the result of the percent inhibition obtained, a straight-line chart will be formed so that there is a linear equation $y = 6.934X + 12.52$ and the value $R^2 = 0.9988$ (figure 2). This equation is used to find IC_{50} value and obtain IC_{50} of vitamin C was 5.40 $\mu\text{g}/\text{mL}$.

4. DISCUSSION

Phytochemical Screening

In the results of research on basil leaf extract during quantitative phytochemical screening test showed that basil leaves (*Ocimum x africanum lour*) compounds such as alkaloids, anthocyanins, cardio glycosides, coumarins, flavonoids, glycosides, phenolic saponins, quinone, steroids, terpenoids, and tannins are in line with research conducted ononamadu et al [38] basil with species *Ocimum canum* containing saponins, phenolics, flavonoids, tannins, carotenoids, triterpenes, glycosides and alkaloids. According to Leila et al [12] *Ocimum basilicum L* contains steroids and Anthocyanins. According to Borah et al [13] *Ocimum Sanctum* contains cardiac glycoside, quinone, and saponins.

Total antioxidant capacity Test

In this study, researchers conducted a total capacity test of antioxidants using DPPH and the ingredients used as a standard comparison is vitamin C (ascorbic acid). The curve of vitamin C has a value of $R^2 = 0.9983$ so that it gets a linear equation $Y = 6.935*X + 12.52$. From the equation, IC₅₀ vitamin C was obtained at 5.40 $\mu\text{g/mL}$. Dpph test curve of basil leaves extract has $R^2 = 0.933$ so that linear equation $Y = 0.1974*X + 15.645$. From the equation, IC₅₀ was obtained at 174.04 $\mu\text{g/mL}$. Based on the results of the DPPH scavenging assay test by measuring IC₅₀ basil leaves extract, IC₅₀ of basil leave was 174.04 $\mu\text{g/mL}$. When compared to IC₅₀ of vitamin C which has a value of 5.40 $\mu\text{g/mL}$, it can be seen that vitamin C has a higher potential to reduce free radicals because the smaller IC₅₀ indicates the higher the total capacity of antioxidant ingredients. Based on Blois [14] criteria, IC₅₀ values of 150-200 $\mu\text{g/mL}$ indicate weak total antioxidant and active capacity [15]. There is little difference with research conducted by Herani et al [15], antioxidant levels in *Ocimum basilicum* leave extract of 141.59 $\mu\text{g/mL}$, this difference is due to different species and different solvents used. Although basil leaves have a lower total antioxidant capacity than vitamin C It must be considered that Vitamin C can induced acid reflux that are not suitable for the people with Gastroesophageal reflux so basil leaves can be the solution for this, beside that basil leave not only contain antioxidant but also contain Linalool and citral which are terpenoid groups that have antioxidants capacity antimicrobials activity and widely used as cosmetic Ingredient [16].




Figure 1 DPPH test of Basil Extract

Vitamin C as comparison standards

The results of absorbance measurement of each concentration of vitamin C and % inhibition using the UV-Vis spectrophotometer tool and created a linear equation curve to obtain the IC₅₀. From the standard curve that is obtained linear equation $Y = 6882*X + 12.62$ and value $R^2 = 0.9988$ obtained $IC_{50} = 5.40 \mu\text{g/mL}$.




Figure 2. Standard Curve of Ascorbic Acid

5. CONCLUSION AND SUGGESTION

Based on the results [36] and discussion on the study of antioxidant capacity test of basil leaf extract, *Ocimum x africanum lour* then obtained conclusions in the below that:

1. Basil leaves have compounds such as alkaloids, anthocyanins, cardio glycosides, coumarins, flavonoids, glycosides, phenolic, saponins, quinone, steroids, terpenoids, and tannins
2. Basil leaves have Antioxidant capacity with its inhibitory concentration 50 (IC₅₀) was 174.04 $\mu\text{g/mL}$, that is categorized as active antioxidant capacity.

More research is needed on other parts besides leaves such as stems and roots and in-vivo research is needed.

ACKNOWLEDGMENTS

The author thanks many colleagues whose research work contributed to this antioxidant research, and we apologize to many author whose interesting work could not be referred to explicitly. It is an honor to acknowledge F. ferdinal for the discussions. Generous research support is provided by Department of Biochemistry and Molecular Biology, Faculty of Medicine, Tarumanagara University

AUTHOR CONTRIBUTIONS

The authors contributed equally to all aspects of the article.

CONFLICT OF INTERESTS

The authors declare that there is no conflict of interest

REFERENCES

[1] E.M. Clarke, E.A. Emerson, Design and synthesis of synchronization skeletons using branching time temporal logic, in: D. Kozen (Eds.), *Workshop on Logics of Programs*, Lecture Notes in Computer Science, vol. 131, Springer, Berlin, Heidelberg, 1981, pp. 52–71. DOI: <https://doi.org/10.1007/BFb0025774>

[2] J.P. Queille, J. Sifakis, Specification and verification of concurrent systems in CESAR, in: M. Dezani-Ciancaglini and U. Montanari (Eds.), *Proceedings of the 5th International Symposium on Programming*, Lecture Notes in Computer Science, vol. 137, Springer, Berlin, Heidelberg, 1982, pp. 337–351. DOI: https://doi.org/10.1007/3-540-11494-7_22

[3] C. Baier, J.-P. Katoen, *Principles of Model Checking*, MIT Press, 2008.

[4] M. Kwiatkowska, G. Norman, D. Parker, Stochastic model checking, in: M. Bernardo, J. Hillston (Eds.), *Proceedings of the Formal Methods for the Design of Computer, Communication and Software Systems: Performance Evaluation (SFM)*, Springer, Berlin, Heidelberg, 2007, pp. 220–270. DOI: https://doi.org/10.1007/978-3-540-72522-0_6

[5] V. Forejt, M. Kwiatkowska, G. Norman, D. Parker, Automated verification techniques for probabilistic systems, in: M. Bernardo, V. Issarny (Eds.), *Proceedings of the Formal Methods for Eternal Networked Software Systems (SFM)*, Springer, Berlin, Heidelberg, 2011, pp. 53–113. DOI: https://doi.org/10.1007/978-3-642-21455-4_3

[6] G.D. Penna, B. Intrigila, I. Melatti, E. Tronci, M.V. Zilli, Bounded probabilistic model checking with the muralpha verifier, in: A.J. Hu, A.K. Martin (Eds.), *Proceedings of the Formal Methods in Computer-Aided Design*, Springer, Berlin, Heidelberg, 2004, pp. 214–229. DOI: https://doi.org/10.1007/978-3-540-30494-4_16

[7] E. Clarke, O. Grumberg, S. Jha, et al., Counterexample-guided abstraction refinement, in: E.A. Emerson, A.P. Sistla (Eds.), *Computer Aided Verification*, Springer, Berlin, Heidelberg, 2000, pp. 154–169. DOI: https://doi.org/10.1007/10722167_15

[8] H. Barringer, R. Kuiper, A. Pnueli, Now you may compose temporal logic specifications, in: *Proceedings of the Sixteenth Annual ACM Symposium on the Theory of Computing (STOC)*, ACM, 1984, pp. 51–63. DOI: <https://doi.org/10.1145/800057.808665>.

[9] A. Pnueli, In transition from global to modular temporal reasoning about programs, in: K.R. Apt (Ed.), *Logics and Models of Concurrent Systems*, Springer, Berlin, Heidelberg, 1984, pp. 123–144. DOI: https://doi.org/10.1007/978-3-642-82453-1_5

[10] B. Meyer, Applying "Design by Contract", *Computer* 25(10) (1992) 40–51. DOI: <https://doi.org/10.1109/2.161279>

[11] S. Bensalem, M. Bogza, A. Legay, T.H. Nguyen, J. Sifakis, R. Yan, Incremental component-based construction and verification using invariants, in: *Proceedings of the Conference on Formal Methods in Computer Aided Design (FMCAD)*, IEEE Press, Piscataway, NJ, 2010, pp. 257–256.

[12] H. Barringer, C.S. Pasareanu, D. Giannakopoulou, Proof rules for automated compositional verification through learning, in: *Proc. of the 2nd International Workshop on Specification and Verification of Component Based Systems*, 2003.

[13] M.G. Bobaru, C.S. Pasareanu, D. Giannakopoulou, Automated assume-guarantee reasoning by abstraction refinement, in: A. Gupta, S. Malik (Eds.), *Proceedings of the Computer Aided Verification*, Springer, Berlin, Heidelberg, 2008, pp. 135–148. DOI: https://doi.org/10.1007/978-3-540-70545-1_14

Basil_Leaf_Extract_Ocimum_x_africanum_Lour

ORIGINALITY REPORT



PRIMARY SOURCES

1	Submitted to University of Suffolk Student Paper	2%
2	Henzinger, T.A.. "On the universal and existential fragments of the @m-calculus", Theoretical Computer Science, 20060328 Publication	1%
3	www.coursehero.com Internet Source	1%
4	Submitted to University of Birmingham Student Paper	1%
5	Submitted to University of Wisconsin-Whitewater Student Paper	1%
6	journal.universitaspahlawan.ac.id Internet Source	1%
7	Maura Lodovici, Elisabetta Bigagli. "Oxidative Stress and Air Pollution Exposure", Journal of Toxicology, 2011 Publication	1%

8 Zhi Yuan Chen, Shao Bin Huang, Li Li Han. "A Fast Approach of Locating Complex System Design Errors", Key Engineering Materials, 2013 1 %
Publication

9 Submitted to Universitas Mulawarman 1 %
Student Paper

10 Bustan, D.. "Applicability of fair simulation", Information and Computation, 20041010 1 %
Publication

11 media.neliti.com 1 %
Internet Source

12 Siufui Hendrawan, Jennifer Lheman, Nuraeni, Ursula Weber, Hans Ulrich Baer. "Hepatocyte and Islet Cell Cotransplantation on Poly-L-Lactide Matrix for the Treatment of Liver Cirrhosis", International Journal of Hepatology, 2020 1 %
Publication

13 Ferreira-Cravo, M.. "Antioxidant responses and reactive oxygen species generation in different body regions of the estuarine polychaeta *Laeonereis acuta* (Nereididae)", Chemosphere, 200701 1 %
Publication

14 Submitted to Ashesi University 1 %
Student Paper

15	Submitted to Bemidji State University Student Paper	1 %
16	Siufui Hendrawan, Epifania Bono, Albert Hutter, Ursula Weber, Jennifer Lheman, Hans U. Baer. " Evaluation of scaffolds coated with collagen as carrier for hepatocytes ", Journal of Biomedical Materials Research Part B: Applied Biomaterials, 2020 Publication	1 %
17	docplayer.net Internet Source	1 %
18	pure.tue.nl Internet Source	1 %
19	d.docksci.com Internet Source	<1 %
20	ijomeh.eu Internet Source	<1 %
21	Buchholz, P.. "Model-checking large structured Markov chains", Journal of Logic and Algebraic Programming, 200305/08 Publication	<1 %
22	Submitted to Higher Education Commission Pakistan Student Paper	<1 %
23	N Tarigan, E Julianti, J Silalahi, H Sinaga. "Promoting sustainability of herbs and spices	<1 %

from Simalungun based on its bioactive compound in traditional food tinuktuk", IOP Conference Series: Earth and Environmental Science, 2023

Publication

24	dspace.univ-ouargla.dz	<1 %
25	Kristina Timonen. "Making meaning of an artwork: promoting expression of thoughts through artwork for patients with eating disorders", <i>Social Semiotics</i> , 2023	<1 %
	Publication	
26	Submitted to University of Oxford	<1 %
	Student Paper	
27	doi.org	<1 %
	Internet Source	
28	repository.ipb.ac.id:8080	<1 %
	Internet Source	
29	Submitted to CSU, San Jose State University	<1 %
	Student Paper	
30	Submitted to HTM (Haridus- ja Teadusministeerium)	<1 %
	Student Paper	
31	Submitted to Bonn International School	<1 %
	Student Paper	

32	Submitted to Krida Wacana Christian University Student Paper	<1 %
33	www.hindawi.com Internet Source	<1 %
34	www.mdpi.com Internet Source	<1 %
35	apps.dtic.mil Internet Source	<1 %
36	repo.itera.ac.id Internet Source	<1 %
37	text-id.123dok.com Internet Source	<1 %
38	www.ncbi.nlm.nih.gov Internet Source	<1 %
39	islamicmarkets.com Internet Source	<1 %
40	Alice Miller. "Symmetry in temporal logic model checking", ACM Computing Surveys, 9/30/2006 Publication	<1 %
41	Sajjad Ali, Muhammad Abdul Basit-Ur-Rahim, Fahim Arif. "Formal verification of internal block diagram of SysML for modeling real-time system", 2015 IEEE/ACIS 16th	<1 %

International Conference on Software Engineering, Artificial Intelligence, Networking and Parallel/Distributed Computing (SNPD), 2015

Publication

Exclude quotes Off
Exclude bibliography Off

Exclude matches Off